

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/68, G01N 33/48	A1	(11) International Publication Number: WO 94/21822
		(43) International Publication Date: 29 September 1994 (29.09.94)

(21) International Application Number: PCT/US94/02938
(22) International Filing Date: 18 March 1994 (18.03.94)
(30) Priority Data:
08/034,738 19 March 1993 (19.03.93) US

(71)(72) Applicant and Inventor: KÖSTER, Hubert [DE/US];
1640 Monument Street, Concord, MA 01742 (US).

(74) Agents: DECONTI, Giulio, A. et al.; Lahive & Cockfield, 60
State Street, Boston, MA 02109 (US).


(81) Designated States: AU, CA, CN, JP, RU, European patent
(AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published
With international search report.

(54) Title: DNA SEQUENCING BY MASS SPECTROMETRY VIA EXONUCLEASE DEGRADATION

(57) Abstract

Methods for determining the sequence of nucleic acids by cleaving the nucleic acid unilaterally from a first end with an exonuclease activity to sequentially release individual nucleotides, identifying each of the sequentially release nucleotides by mass spectrometry, and determining the sequence of the nucleic acid from the identified nucleotides are disclosed. The method is amenable to multiplexing for simultaneously determining more than one nucleic acid sequence.



Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

GB	United Kingdom
GE	Georgia
GN	Guinea
GR	Greece
HU	Hungary
IE	Ireland
IT	Italy
JP	Japan
KE	Kenya
KG	Kyrgyzstan
KP	Democratic People's Republic of Korea
KR	Republic of Korea
KZ	Kazakhstan
LI	Liechtenstein
LK	Sri Lanka
LU	Luxembourg
LV	Latvia
MC	Monaco
MD	Moldova

MR	Mauritania
MW	Malawi
NE	Niger
NL	Netherlands
NO	Norway
NZ	New Zealand
PL	Poland
PT	Portugal
RO	Romania
RU	Russian Federation
SD	Sudan
SE	Sweden
SI	Slovenia
SK	Slovakia
SN	Senegal
TD	Chad
TG	Togo
TJ	Tajikistan
TT	Trinidad and Tobago

*DNA Sequencing By Mass Spectrometry Via
Exonuclease Degradation*

5 **Background Of The Invention**

The fundamental role that determining DNA sequences has for the life sciences is evident. Its importance in the human genome project has been discussed and published widely [e.g. J.E. Bishop and M. Waldholz, 1991, *Genome. The Story of the Most Astonishing Scientific Adventure of Our Time - The Attempt to Map All Genes in the*

10 *Human Body*, Simon & Schuster, New York].

The current state-of-the-art in DNA sequencing is summarized in recent review articles [e.g. B. Barrell, *The FASEB Journal*, 5, 40 (1991); G.L. Trainor, *Anal. Chem.* 62, 418 (1990), and references cited therein]. The most widely used DNA sequencing chemistry is the enzymatic chain termination method [F. Sanger et al., *Proc. Natl. Acad. Sci.*

15 USA, 74, 5463 (1977)] which has been adopted for several different sequencing strategies. The sequencing reactions are either performed in solution with the use of different DNA polymerases, such as the thermophilic Taq DNA polymerase [M.A. Innes, *Proc. Natl. Acad. Sci. USA*, 85, 9436 (1988)] or specially modified T7 DNA polymerase ("SEQUENASE") [S. Tabor and C.C. Richardson, *Proc. Natl. Acad. Sci. USA*, 84, 4767 (1987)], or in conjunction

20 with the use of polymer supports. See for example S. Stahl et al., *Nucleic Acids Res.*, 16, 3025 (1988); M. Uhlen, PCT Application WO 89/09282; Cocuzza et al., PCT Application WO 91/11533; and Jones et al., PCT Application WO 92/03575, incorporated by reference herein.

A central, but at the same time limiting, part of almost all sequencing

25 strategies used today is the separation of the base-specifically terminated nested fragment families by polyacrylamide gel electrophoresis (PAGE). This method is time-consuming and error prone and can result in ambiguous sequence determinations. As a consequence of the use of PAGE, highly experienced personnel are often required for the interpretation of the sequence ladders obtained by PAGE in order to get reliable results. Automatic sequence

30 readers very often are unable to handle artifacts such as "smiling", compressions, faint ghost bands, etc. This is true for the standard detection methods employing radioactive labeling such as ³²P, ³³P or ³⁵S, as well as for the so-called Automatic DNA Sequencers (e.g. Applied Biosystems, Millipore, DuPont, Pharmacia) using fluorescent dyes for the detection of the sequencing bands.

35 Apart from the time factor the biggest limitation of all methods involving PAGE as an integral part, however, is the generation of reliable sequence information, and the transformation insert of this information into a computer format to facilitate subsequent

With standard Sanger sequencing 200 to 500 bases of unconfirmed sequence information could be obtained in about 24 hours; with automatic DNA sequencers this number could be multiplied by approximately a factor of 10 to 20 due to processing several samples simultaneously. A further increase in throughput could be achieved by employing
5 multiplex DNA sequencing [G. Church et al., Science, **240**, 185-188 (1988); Köster et al.,
Nucleic Acids Res. Symposium Ser. No. 24, 318-21 (1991)] in which, by using a unique tag
sequence, several sequencing ladders could be detected one after the other from the same
PAGE after blotting, UV-crosslinking to a membrane, and hybridizations with specific
10 complementary tag probes. However, this approach is still very laborious, often requires
highly skilled personnel and can be hampered by the use of PAGE as a key element of the
whole process.

A large scale sequencing project often starts with either a cDNA or genomic library of large DNA fragments inserted in suitable cloning vectors such as cosmid, plasmid (e.g. pUC), phagemid (e.g. pEMBL, pGEM) or single-stranded phage (e.g. M13) vectors [T.
15 Maniatis, E.F. Fritsch and J. Sambrook (1982) Molecular Cloning. A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.; Methods in Enzymology, Vol.
101 (1983), Recombinant DNA, Part C; Vol. 153 (1987), Recombinant DNA, Part D; Vol.
154 (1987), Recombinant DNA, Part E; Vol. 155 (1987), Recombinant DNA, Part F and Vol.
152 (1987), Guide to Molecular Cloning Techniques, Academic Press, New York]. Since
20 large DNA fragments currently cannot be sequenced directly in one run because the Sanger
sequencing chemistry allows only about 200 to 500 bases to be read at a time, the long DNA
fragments have to be cut into shorter pieces which are separately sequenced. In one approach
this is done in a fully random manner by using, for example, unspecific DNase I digestion,
frequently cutting restriction enzymes, or sonification, and sorting by electrophoresis on
25 agarose gels [Methods in Enzymology, *supra*]. However, this method is time-consuming and
often not economical as several sequences are sequenced many times until a contiguous DNA
sequence is obtained. Very often the expenditure of work to close the gaps of the total
sequence is enormous. Consequently, it is desirable to have a method which allows
sequencing of a long DNA fragment in a non-random, i.e., direct, way from one end through
30 to the other. Several strategies have been proposed to achieve this [Methods of Enzymology,
supra; S. Henikoff, Gene, **28**, 351-59 (1984); S. Henikoff, et al. US Patent No. 4,843,003; and
PCT Application WO 91/12341]. However, none of the currently available sequencing
methods provide an acceptable method of sequencing megabase DNA sequences in either a
timely or economical manner. The main reason for this stems from the use of PAGE as a
35 central and key element of the overall process.

In PAGE, under denaturing conditions the nested families of terminated DNA fragments are separated by the different mobilities of DNA chains of different length. A

the mobility [R. Frank and H. Köster, Nucleic Acids Res., **6**, 2069 (1979)]. PAGE therefore is not only a very slow, but also an unreliable method for the determination of molecular weights, as DNA fragments of the same length but different sequence/base composition could have different mobilities. Likewise, DNA sequences which have the same mobility could have different sequence/base compositions.

The most reliable way for the determination of the sequence/base composition of a given DNA fragment would therefore be to correlate the sequence with its molecular weight. Mass spectrometry is capable of doing this. The enormous advantage of mass spectrometry compared to the above mentioned methods is the speed, which is in the range of seconds per analysis, and the accuracy of mass determination, as well as the possibility to directly read the collected mass data into a computer. The application of mass spectrometry for DNA sequencing has been investigated by several groups [e.g. Methods in Enzymology, Vol. 193: Mass Spectrometry, (J.A. McCloskey, editor), 1990, Academic Press, New York; K.H. Schramm Biomedical Applications of Mass Spectrometry, **34**, 203-287 (1990); P.F. Crain Mass Spectrometry Reviews, **9**, 505 (1990)].

Most of the attempts to use mass spectrometry to sequence DNA have used stable isotopes for base-specific labeling, as for instance the four sulfur isotopes ^{32}S , ^{33}S , ^{34}S and ^{36}S . See, for example, Brennan et al., PCT Application WO 89/12694, R.L. Mills United States Patent No. 5,064,754, United States Patent No. 5,002,868, Jacobson et al.; Hean European Patent Application No. A1 0360676. Most of these methods employ the Sanger sequencing chemistry and - which jeopardizes to some extent the advantages of mass spectrometry - polyacrylamide gel electrophoresis with some variations such as capillary zone electrophoresis (CZE) to separate the nested terminated DNA fragments prior to mass spectrometric analysis.

One advantage of PAGE is the property of being a parallel method, i.e., several samples could be analyzed simultaneously (though this is not true for CZE which is a serial method), whereas mass spectrometry allows, in general, only a serial handling of the samples. In a US Patent Application No.08/001,323, mass spectrometric DNA sequencing is proposed without the use of PAGE, employing desorption/ionization techniques applicable to larger biopolymers such as electrospray (ES) [J.B. Fenn et al., J. Phys. Chem., **88**, 4451-59 (1984); Fenn et al., PCT Application No. WO 90/14148; and B. Ardrey, Spectroscopy Europe, **4**, 10-18 (1992)] and matrix assisted laser desorption/ionization (MALDI) mass spectrometry [F. Hillenkamp et al., Laser Desorption Mass Spectrometry, Part I: Mechanisms and Techniques and Part II: Performance and Application of MALDI of Large Biomolecules, in Mass Spectrometry in the Biological Sciences: A Tutorial (M.L. Gross, editor), 165-197 (1992), Kluwer Academic Publishers, The Netherlands] which can facilitate determination of DNA sequences by direct measurement of the molecular masses in the mixture of base-

analysis typical for current mass spectrometric methods can be changed to a parallel mode [H. Köster, US Patent Application No. 08/001,323, *supra*].

MALDI and ES mass spectrometry are in some aspects complementary techniques. While ES, using an atmospheric pressure ionization interface (API), can accommodate continuous flow streams from high-performance liquid chromatographs (HPLC) [K.B. Tomer, et al. *Biological Mass Spectrometry*, 20, 783-88 (1991)] and capillary zone electrophoresis (CZE) [R.D. Smith et al., *Anal. Chem.*, 60, 436-41 (1988)] this is currently not available for MALDI mass spectrometry. On the other hand, MALDI mass spectrometry is less sensitive towards buffer salts and other low molecular weight components in the analysis of larger molecules with a TOF mass analyzer [Hillenkamp et al. (1992), *supra*]; in contrast ES is very sensitive towards by-products of low volatility. While the high mass range in ES mass spectrometry is accessible through the formation of multiply charged molecular ions, this is achieved in MALDI mass spectrometry by applying a time-of-flight (TOF) mass analyzer and the assistance of an appropriate matrix to volatilize the biomolecules. Similar to ES, a thermospray interface has been used to couple HPLC on-line with a mass analyzer. Nucleosides originating from enzymatic hydrolysates have been analyzed using such a configuration [C.G. Edmonds et al. *Nucleic Acids Res.*, 13, 8197-8206 (1985)]. However, Edmonds et al. does not disclose a method for nucleic acid sequencing.

A complementary and completely different approach to determine the DNA sequence of a long DNA fragment would be to progressively degrade the DNA strand using exonucleases from one side - nucleotide by nucleotide. This method has been proposed by Jett et al. See J.H. Jett et al. *J. Biomolecular Structure & Dynamics*, 7, 301-309 (1989); and J.H. Jett et al. PCT Application No. WO 89/03432. A single molecule of a DNA or RNA fragment is suspended in a moving flow stream and contacted with an exonuclease which cleaves off one nucleotide after the other. Detection of the released nucleotides is accomplished by specifically labeling the four nucleotides with four different fluorescent dyes and involving laser induced flow cytometric techniques.

However strategies which use a stepwise enzymatic degradation process can suffer from the problem that this process is difficult to synchronize, i.e., the enzymatic reaction soon comes out of phase. Jett et al., *supra*, have attempted to address this problem by degrading just one single DNA or RNA molecule by an exonuclease. However, this approach is very hard, as handling a single molecule, keeping it in a moving flow stream, and achieving a sensitivity of detection which clearly identifies one single nucleotide are only some of the very difficult technical problems to be solved. In addition, in using fluorescent tags, the physical detection process for a fluorescent signal involves a time factor difficult to control and the necessity to employ excitation by lasers can cause photo-bleaching of the fluorescent signal.

systems suitable for high-speed DNA sequencing a prerequisite to tackle the human genome (and other genome) sequencing projects.

Summary Of The Invention

5 In contrast to most sequencing strategies, the process of this invention does not use the Sanger sequencing chemistry, polyacrylamide gel electrophoresis or radioactive, fluorescent or chemiluminescent detection. Instead the process of the invention adopts a direct sequencing approach, beginning with large DNA fragments cloned into conventional cloning vectors, and based on mass spectrometric detection. To achieve this, the DNA is by
10 means of protection, specificity of enzymatic activity, or immobilization, unilaterally degraded in a stepwise manner via exonuclease digestion and the nucleotides or derivatives detected by mass spectrometry.

 Prior to this enzymatic degradation, sets of ordered deletions can be created which span the whole sequence of the cloned DNA fragment. In this manner, mass-modified
15 nucleotides can be incorporated using a combination of exonuclease and DNA/RNA polymerase. This enables either multiplex mass spectrometric detection, or modulation of the activity of the exonuclease so as to synchronize the degradative process. In another embodiment of the invention the phasing problem can be resolved by continuously applying small quantities of the enzymatic reaction mixture onto a moving belt with adjustable speed
20 for mass spectrometric detection. In yet another embodiment of the invention, the throughput could be further increased by applying reaction mixtures from different reactors simultaneously onto the moving belt. In this case the different sets of sequencing reactions can be identified by specific mass-modifying labels attached to the four nucleotides. Two-dimensional multiplexing can further increase the throughput of exonuclease mediated mass
25 spectrometric sequencing as being described in this invention.

Brief Description Of The Drawings

The accompanying drawings form a part of the specifications and serve the purpose of providing examples illustrating certain embodiments of the present invention. Together with the description they help to explain the principles of the invention without
5 limiting the scope of it.

FIGURE 1 illustrates a process of exonuclease sequencing beginning with a single-stranded nucleic acid.

FIGURE 2 illustrates a process similar to FIGURE 1, however, starting with a target nucleic acid inserted into a double-stranded vector.

10 FIGURE 3 illustrates a method for introducing mass-modified nucleotides into a target nucleic acid sequence multiplexing mass spectrometry.

FIGURE 4A and 4B illustrate methods for introducing mass-modified nucleotides into a target nucleic acid sequence multiplexing mass spectrometry.

15 FIGURE 5 shows positions within a nucleic acid molecule which can be modified for the introduction of discriminating mass increments or modulation of exonuclease activity.

FIGURE 6 illustrates various structures of modified nucleoside triphosphates useful for the enzymatic incorporation of mass-modified nucleotides into the DNA or RNA to be sequenced.

20 FIGURE 7 shows some possible functional groups (R) useful to either mass-modification of nucleotides in discrete increments for differentiation by mass spectrometry and/or to modulate the enzymatic activity of an exonuclease.

FIGURE 8 illustrates some linking groups X for the attachment of the mass modifying functionality R to nucleosides.

25 FIGURE 9 is a schematic drawing of a sequencing reactor system.

FIGURE 10 is a graphical representation of idealized output signals following the time-course of the stepwise mass spectrometric detection of the exonucleolytically released nucleotides.

30 FIGURE 11 illustrates specific labels introduced by mass modification to facilitate multiplex exonuclease mass spectrometric sequencing.

FIGURE 12 is a schematic drawing of a moving belt apparatus for delivering single or multiple tracks of exonuclease samples for laser induced mass spectrometric sequence determination in conjunction with the sequencing reactor of FIGURE 9.

35 FIGURE 13 is a schematic representation of individually labeled signal tracks employed in multiplex exonuclease mediated mass spectrometric sequencing.

FIGURES 14A and 14B illustrate a method for double-stranded exonuclease sequencing for mass spectrometric sequence determination.

Detailed Description Of The Invention

The starting point for the process of the invention can be, for example, DNA cloned from either a genomic or cDNA library, or a piece of DNA isolated and amplified by polymerase chain reaction (PCR) which contains a DNA fragment of unknown sequence.

5 Libraries can be obtained, for instance, by following standard procedures and cloning vectors [Maniatis, Fritsch and Sambrook (1982), *supra*; Methods in Enzymology, Vol. 101 (1983) and Vol. 152-155 (1987), *supra*]. Appropriate cloning vectors are also commercially available. As will be apparent, the invention is not limited to the use of any specific vector constructs and cloning procedures, but can be applied to any given DNA fragment whether
10 obtained, for instance, by cloning in any vector or by the Polymerase Chain Reaction (PCR). The unknown DNA sequence can be obtained in either double-stranded form using standard PCR or in a single-stranded form employing asymmetric PCR [PCR Technology: Principles and Applications for DNA Amplification (Erlich, editor), M. Stockton Press, New York (1989)].

15 For those skilled in the art it is clear that both DNA and RNA can be exonucleolytically degraded from either the 5' or 3' end depending upon the choice of exonuclease. Similarly the sequence of an unknown DNA molecule can be determined directly by exonuclease digestion, or alternatively, the DNA fragment of unknown sequence can be transcribed first into a complementary RNA copy which is subsequently
20 exonucleolytically degraded to provide the RNA sequence. Appropriate vectors, such as the pGEM (Promega) vectors, are useful in the present as they have specific promoters for either the SP6 or T7 DNA dependent RNA polymerases flanking the multiple cloning site. This feature allows transcription of both unknown DNA strands into complementary RNA for subsequent exonuclease sequencing. Furthermore, these vectors, belonging to the class of
25 phagemid vectors, provide means to generate single stranded DNA of the unknown double stranded DNA. Thus, by using two vectors which differ only in the orientation of the fl origin of replication, both strands of the unknown DNA sequence can be obtained in a single stranded form and utilized for subsequent exonuclease sequencing. The scope of the invention is also not limited by the choice of restriction sites. There is, however, a preference
30 for rare cutting restriction sites to keep the unknown DNA fragment unfragmented during the manipulations in preparation for exonuclease sequencing. The following description and the FIGURES do serve the purpose by way of examples to illustrate the principles of the invention. Variations which are within the scope of the invention will be obvious to those skilled in the art. Various mass spectrometric configurations are within the scope of the
35 invention: For desorption/ionization e.g. fast atomic bombardment (FAB), plasma desorption (PD), thermospray (TS) and preferentially electrospray (ES) and laser desorption with and without an appropriate matrix (LD or

- (i) Preparation of unknown nucleic acid sequence for exonuclease sequencing:

FIGURE 1 describes the process for a single stranded DNA insert ("Target DNA") of a single stranded circular cloning vector. The boundaries of the target DNA are designated A and B. The target DNA, as illustrated in FIGURE 1, has been cloned into the Not I site of a vector. A synthetic oligodeoxynucleotide [N.D. Sinha, J. Biernat, J. McManus and H. Köster, *Nucleic Acids Res.*, **12**, 4539 (1984)] which will restore the Not I site to double-strandedness and which is complementary to the vector sequence flanking the A boundary of the insert DNA is hybridized to that site and cleaved by Not I restriction endonuclease. The two pieces of the synthetic oligodeoxynucleotide can then be removed by molecular sieving, membrane filtration, precipitation, or other standard procedures.

FIGURE 1 also illustrates a set of ordered deletions (t^0 , t^1 , t^2 , t^3) which can be obtained by the time-limited action of an exonuclease, e.g. T4 DNA polymerase, in the absence of dNTPs. The set of deletions can be immobilized on a solid support Tr (Tr^0 , Tr^1 , Tr^2 , Tr^3), or alternatively, the set of ordered deletions can be obtained in a heterogeneous reaction by treating the solid support Tr^0 containing the complete target DNA sequence with an exonuclease in a time-limited manner. In the instance where the 3' termini of each time point are too heterogeneous (i.e., "fuzzy") to be analyzed directly by exonuclease mediated mass spectrometric sequencing, circularization of the template and a cloning step can be performed prior to this sequencing process with single transformed colonies selected.

A single stranded linear DNA fragment carrying the unknown sequence with its A boundary at the 3' end can be directly sequenced by an 3' exonuclease in an apparatus described below and schematically depicted in FIGURE 9, provided that the exonuclease is immobilized within the reactor on, for example, beads, on a frit, on a membrane located on top of the frit or on the glass walls of a capillary or entrapped in a gel matrix or simply by a semipermeable membrane which keeps the exonuclease in the reactor while the linear DNA fragment is circulating through a loop.

At time intervals, or alternatively as a continuous stream, the reaction mixture containing the buffer and the released nucleotides is fed to the mass spectrometer for mass determination and nucleotide identification. In another embodiment, the stream containing the nucleotides released by exonuclease action can be passed through a second reactor or series of reactors which cause the released nucleotide to be modified. For example, the second reactor can contain an immobilized alkaline phosphatase and the nucleotides passing therethrough are transformed to nucleosides prior to feeding into the mass spectrometer. Other mass-modifications are described below.

In general, when it is the released nucleotide (or ribonucleotide) which is

synthesis can also be used to modify the released nucleotide just prior to mass spectrometric analysis. For instance, the amino function of adenine, guanine or cytosine can be modified by acylation. The amino acyl function can be, by way of illustration, an acetyl, benzoyl, isobutyryl or anisoyl group. Benzoylchloride, in the presence of pyridine, can acylate the adenine amino group, as well as the deoxyribose (or ribose) hydroxyl groups. As the glycosidic linkage is more susceptible to hydrolysis, the sugar moiety can be selectively deacylated if the acyl reaction was not efficient at those sites (i.e., heterogeneity in molecular weight arising from incomplete acylation of the sugar). The sugar moiety itself can be the target of the mass-modifying chemistry. For example, the sugar moieties can be acylated, tritylated, monomethoxytritylated, etc. Other chemistries for mass-modifying the released nucleotides (or ribonucleotides) will be apparent to those skilled in the art.

In another embodiment the linear single stranded DNA fragment can be anchored to a solid support. This can be achieved, for example, by covalent attachment to a functional group on the solid support, such as through a specific oligonucleotide sequence which involves a spacer of sufficient length for the ligase to react and which is covalently attached via its 5' end to the support (FIGURE 1). A splint oligonucleotide with a sequence complementary in part to the solid bound oligonucleotide and to the 5' end of the linearized single stranded vector DNA allows covalent attachment of the DNA to be sequenced to the solid support. After annealing, ligation (i.e., with T4 DNA ligase) covalently links the solid bound oligonucleotide and the DNA to be sequenced. The splint oligonucleotide can be subsequently removed by a temperature jump and/or NaOH treatment, or washed off the support using other standard procedures. The solid support with the linear DNA example is transferred to the reactor (FIGURE 9) and contacted with an exonuclease in solution. As illustrated, where the 3' end of the unknown DNA fragment is exposed (ie unprotected), a 3' exonuclease is employed. The released nucleotides, or modified nucleotides if intermediately contacted with a modifying agent such as alkaline phosphatase, are identified by mass spectrometry as described above. Other linking groups are described herein, and still others will be apparent to those skilled in the art from the invention described here.

The solid supports can be of a variety of materials and shapes, such as beads of silica gel, controlled pore glass, cellulose, polyacrylamide, sepharose, sephadex, agarose, polystyrene and other polymers, or membranes of polyethylene, polyvinylidendifluoride (PVDF) and the like. The solid supports can also be capillaries, as well as frits from glass or polymers. Various 3' exonucleases can be used, such as phosphodiesterase from snake venom, Exonuclease VII from *E. coli*, Bal 31 exonuclease and the 3'-5' exonuclease activity of some DNA polymerases exerted in the absence of dNTPs, as for example T4 DNA polymerase.

In using a phagemid vector with an inverted fl origin of replication, the R

1990; 1: 277-287) and the effect of endonuclease, by phagizing

oligodeoxynucleotide and splint oligonucleotide. As another embodiment of this invention, the hybridizing oligonucleotide can also be designed to bind a promoter site upstream of the A boundary and by doing so restoring the doublestranded promoter DNA. Directly, or with a short initiator oligonucleotide carrying an attachment functionality at the 5' end, transcription
5 can be initiated with the appropriate specific DNA dependent RNA polymerase [Methods in Enzymology, Vol. 185, Gene Expression Technology (1990); J.F. Milligan, D.R. Groebe, G.W. Witherell and O.C. Uhlenbeck, Nucleic Acids Res., **15**, 8783-98 (1987); C. Pitulle, R.G. Kleineidam, B. Sproat and G. Krupp, Gene, **112**, 101-105 (1992) and H. Köster US Patent Application No. 08/001,323, supra]. The RNA transcript can be transferred to the
10 reactor (FIGURE 9) and contacted with an immobilized or otherwise contained exonuclease, or immobilized via the 5' functionality of the initiator oligonucleotide incorporated in the RNA transcript to a solid support and then contacted with an exonuclease in solution.

Depending on the length of the DNA insert (i.e., number of nucleotides between boundary A and B in FIGURE 1) the mass spectrometric exonuclease sequencing
15 process can allow the complete sequence from A to B to be determined in one run. Alternatively, prior to exonuclease sequencing, a set of ordered deletions can be prepared according to standard procedures [e.g. Methods in Enzymology, Vol. 101 (1983) and Vol. 152-155 (1987); R.M.K. Dale et al., Plasmid, **13**, 31-40 (1985)], such that, in FIGURE 1 the steps Tr⁰ to Tr³ can represent either different time values of the mass spectrometric
20 exonuclease sequencing reaction from immobilized DNA fragments or different starting points for the exonuclease DNA/RNA mass spectrometric sequencing process. In either case the principle of the invention described provides a process by which the total sequence of the insert can be determined.

In another embodiment of the invention the unknown DNA sequence (target
25 DNA) is inserted into a double-stranded cloning vector (FIGURE 2) or obtained in double-stranded form, as for example by a PCR (polymerase chain reaction) process [PCR Technology, (1989) supra]. The DNA to be sequenced is inserted into a cloning vector, such as ligated into the Not I site as illustrated in Figure 2. Adjacent to the A boundary there can be located another cutting restriction endonuclease site, such as an Asc I endonuclease
30 cleavage site. The double-stranded circular molecule can be linearized by treatment with Asc I endonuclease and ligated to a solid support using a splint oligodeoxynucleotide (and ligase) as described above which restores the Asc I restriction site (Tr⁰ds and Tr⁰'ds). The strand which is not immobilized can be removed by subjecting the double stranded DNA to standard denaturing conditions and washing, thereby generating single-stranded DNAs immobilized to
35 the solid support (Tr⁰ss and Tr⁰'ss). Since the unknown double-stranded DNA sequence can be ligated in either orientation to the support there can exist two non-identical 3' termini (+ and - strand) immobilized, which can result in ambiguous sequencing data. The immobilized fragment which carries the

oligodeoxynucleotide complementary to the 3' end of the strand to be protected. As there can only be hybridization at one 3' terminus, i.e., to the wrong single-stranded DNA with (-) strand information (Tr^{0ss}), some alpha-thio dNTP's can be incorporated into the immobilized (-) strand via treatment with a DNA polymerase and completely protect that strand from
5 exonucleolytic degradation [P.M.J. Burgers and F. Eckstein, Biochemistry, **18**, 592 (1979); S. Labeit, H. Lehrach and R.S. Goody, DNA, **5**, 173 (1986); S. Labeit, H. Lehrach and R.S. Goody in Methods in Enzymology, Vol. 155, page 166 (1987), supra]. If desired, after incorporation of exonuclease resistant nucleotides, the oligonucleotide primer may be removed by a washing step under standard denaturing conditions. The immobilized single-
10 stranded DNAs are transferred to the sequencing reactor (FIGURE 9) and the sample with the unknown sequence at the 3' end is degraded by an exonuclease in a stepwise manner. The liberated nucleotides, or optionally modified nucleotides, are continuously fed into the mass spectrometer to elucidate the sequence.

As above, where the inserted DNA is too long for determining the complete
15 sequence information between the boundaries A and B (FIGURE 2) in one run of exonuclease mass spectrometric sequencing, a series of overlapping ordered deletions can be constructed according to standard procedures, e.g. utilizing the restriction site RIII producing 3' sticky ends inert towards exonuclease III digestion [Methods in Enzymology, Vol 152-155 (1987) and S. Henikoff, Gene, (1984), supra]. When required, the deletion mutants can be
20 recircularized and used to transform host cells following standard procedures. Single colonies of the transformed host cells are selected, further proliferated, and the deletion mutant isolated and immobilized as single-stranded DNAs, similar to the process described above and subsequently analyzed by exonuclease mass spectrometric sequencing. Alternatively, the immobilized full length single stranded DNA (Tr^{0ss}) can be treated in
25 time-limited reactions with an exonuclease such as T4 DNA polymerase in the absence of NTPs, to create a set of immobilized ordered deletions for subsequent exonuclease mass spectrometric sequencing. However, in case the 3' termini are too heterogeneous for direct mass spectrometric exonuclease sequencing an intermediate cloning step can be included. In yet another embodiment, the exonuclease mediated sequencing can be performed by
30 providing the single-stranded (ss) nucleic acid fragment in solution. This can be achieved by treating the solid support, e.g. Tr^{0ss} , with an oligonucleotide complementary to the unique Asc I site. After hybridization this site is now double-stranded (ds) and susceptible to Asc I endonuclease cleavage and release of the single-stranded fragment.

If a cloning vector such as one of the pGEM family (Promega Corp.) is used,
35 both strands of the double-stranded target DNA can be transcribed separately dependant upon which of the specific promoters flanking the insertion site is used (located next to the A or B boundary of the insert, FIGURE 2) and the corresponding specific DNA dependent RNA

Transcription of the DNA dependent RNA is followed by mass spectrometric exonuclease

sequencing using an immobilized or entrapped exonuclease. In an alternate embodiment, the transcription process is initiated via initiator oligonucleotides with a 5' functionality allowing the subsequent immobilization of the RNA transcripts to a solid support [H. Köster, US Patent Application No. 08/001,323, *supra*]; in this case the mass spectrometric sequencing
5 can be performed within the sequencing reactor using an exonuclease in solution. The stepwise liberated ribonucleotides, or modified ribonucleotides (i.e., ribonucleosides generated by passing through a reactor containing immobilized alkaline phosphatase), are fed to the mass spectrometer for sequence determination.

10

(ii) Introduction of mass-modified nucleotides for multiplex exonuclease sequencing:

Since standard mass spectrometry is a serial process, the throughput can be
15 limited. However, in the present invention the throughput can be considerably increased by the introduction of mass-modified nucleotides into the DNA or RNA to be sequenced allowing for a parallel analysis of several nucleic acid sequences simultaneously by mass spectrometry. See H. Köster, US Patent Application No. 08/001,323, *supra*. Low molecular weight nucleic acid components, such as unmodified or mass modified
20 nucleotides/nucleosides, can be analyzed simultaneously by multiplex mass spectrometry.

Mass-modified nucleotides can be incorporated by way of mass modified nucleoside triphosphates precursors using various methods. For example, one can begin with the insert of the target DNA sequence in a single-stranded cloning vector by having a "primer" and a "stopper" oligonucleotide bound to the complementary vector sequences
25 located at the A and B boundary of the insert DNA respectively (FIGURE 3) and a template directed DNA polymerase, preferentially one lacking the 3'- 5' and 5'- 3' exonuclease activity such as Sequenase, version 2.0 (US Biochemicals, derived from T7 DNA polymerase), Taq DNA polymerase or AMV reverse transcriptase. In the illustrative embodiment, the unknown DNA sequence has been inserted in a restriction endonuclease site such as Not I. Adjacent to the A boundary another restriction endonuclease site, such as the Asc I site, can
30 be located within the primer binding site such that the partly double-stranded circular DNA can be cleaved at the unique Asc I site and the mass-modified (-) strand (t^3 in FIGURE 3) isolated by standard procedures (i.e., membrane filtration, molecular sieving, PAGE or agarose gel electrophoresis) and, if desired, coupled to a solid support via a splint
35 oligonucleotide restoring the Asc I site in double-stranded form for ligation by T4 DNA ligase (FIGURE 3). After having removed the splint oligonucleotide the immobilized single-stranded DNA fragment with its B' boundary at the 3' end (i.e., Tr^3) is ready for exonuclease

primer will not hybridize with its 5' terminal sequence to the vector as is shown in FIGURE 3. it will nevertheless allow the covalent attachment of the single-stranded mass-modified DNA to the solid support using the same splint oligonucleotide as described above. In yet another embodiment the primer can carry a non-restriction site sequence information at its 5' end, which may or may not be complementary to the opposite vector sequence, but is complementary to a specific splint oligodeoxynucleotide which allows the covalent attachment to the solid support. The latter two procedures do not require cleavage with a restriction endonuclease and separation of the strands.

The reaction mixture obtained after enzymatic synthesis of the mass-modified (-) strand can be directly joined to the solid support and the circular vector DNA and the stopper oligonucleotide can be removed under denaturing conditions. In yet another embodiment, the generation of a set of ordered deletions of the target DNA sequence information and the incorporation of mass modified nucleotides can be combined by terminating the DNA polymerase reaction at different time intervals (i.e., t^0 , t^1 , t^2 , t^3 , FIGURE 3) to generate a ladder of mass-modified (-) strands. In case the 3' termini of each time point are too heterogeneous for mass spectrometric exonuclease sequencing a circularization and cloning step as described above can be included.

As illustrated in FIGURES 14A and 14B, both the (+) and (-) strand can be exonuclease sequenced simultaneously. Incorporation of mass-modified nucleotides in to one of the (+) or (-) strands can be carried out as described above. In the illustrative embodiment, both the (+) and (-) strands are ligated to solid supports and exonucleased simultaneously. The presence of the mass-modified nucleotides in the (-) strand can allow for differentiation of mass spectrometric signals arising from nucleotides released from both strands. Where exonuclease sequencing can proceed essentially between the A and B boundaries in one pass, the sequence of the (-) strand can be inverted and aligned with the complimentary sequence. An advantage to this approach is the identification of ambiguous sequencing data (i.e., base pair mismatches arising from error of sequencing one of the strands). Alternatively, the full sequence can be obtained from partial exonuclease sequencing of both the (+) and (-) strands provided that sequencing proceeds to an overlapping point on each strand. By searching for the complementary overlapping region of each sequence fragment and aligning the two sequence fragments, the sequence of the remainder of one or both of the strands can be "reconstructed" based on the known sequence of the other. This later example provides a means of sequencing in "one pass" a much larger DNA fragment then would be possible by exonuclease sequencing only one strand.

In using vectors of the phagemid type (e.g. pGEM family, Promega Corp.) both strands of the unknown DNA fragment can be mass-modified by using just the vector which carries the fl origin of replication in the opposite direction

transcription promoter regions flanking the insertion site, restoring the double-stranded promoter site by complementary oligonucleotides [Methods in Enzymology, Vol. 185, (1990); Uhlenbeck et al., Nucleic Acids Res., (1987), supra] and transcribing with appropriate RNA polymerases in the presence of mass-modified ribonucleoside triphosphates. As above,

5 the mass-modified RNA can be directly transferred to the sequencing reactor for mass spectrometric sequencing using an immobilized or entrapped exonuclease. In another embodiment the transcription can be initiated with initiator oligonucleotides [Krupp et al., Gene, (1992), supra] carrying a 5' functionality for subsequent attachment of the mass-modified RNAs to a solid support. In the later instance, the immobilized mass-modified

10 RNAs can be contacted in the sequencing reactor (FIGURE 9) with an exonuclease in solution for mass spectrometric sequencing.

The mass-modification of the immobilized strand starting with the unknown DNA insert in a double-stranded vector (FIGURE 4A) can be introduced starting with a situation similar to Tr⁰ds in FIGURE 2. However, a 5' phosphorylated exonuclease III

15 resistant splint oligonucleotide (i.e., 2',3' dideoxy) is ligated to the (-) strand allowing a unilateral digestion of the (+) strand with exonuclease III (FIGURE 4A). The mass-modifications are then introduced by a filling-in reaction using template dependent DNA polymerases such as Sequenase, version 2.0 (US Biochemicals), Taq DNA polymerase or AMV reverse transcriptase and appropriate mass-modified dNTPs. In another embodiment

20 one can start with a situation similar to Tr⁰ss in FIGURE 2 and by using a (-) primer designed to bind outside the A boundary at the 3' end of the (+) strand, synthesize a mass-modified (-) strand employing mass-modified dNTPs and a DNA dependent DNA polymerase as described above. In one embodiment, there can be a short stretch of sequence between the Not I and the Asc I site to allow this primer to hybridize effectively. This

25 approach can also be carried out by generating a mass modified (+) strand starting with Tr⁰ss (FIGURE 4B). The newly synthesized (+) strand can be isolated from the (-) strand solid support, such as by denaturation, and immobilized via the 5' sequence information of the primer and a splint oligonucleotide which is in part complementary to this and to an oligonucleotide sequence already attached to another solid support (FIGURE 4B). After

30 ligation (i.e., with T4 ligase) the splint oligonucleotide is removed and the immobilized mass-modified single-stranded (+) DNA is transferred to the sequencing reactor (FIGURE 9) and contacted with an exonuclease, such as T4 DNA polymerase in solution, for mass spectrometric sequence determination via the released mass-modified nucleotides.

In accordance with this invention, the mass-modifying functionality can be

35 located at different positions within the nucleotide moiety (FIGURE 5 and 6). See also H. Köster, US Patent Application No. 08/001,323, for further examples and synthesis chemistries. For instance, the mass-modifying functionality can be located at the heterocyclic

(M1). Modifications in these positions do not interfere with Watson-Crick specific base-pairing necessary for the enzymatic incorporation into the mass-modified nucleic acids (DNA/RNA) with high accuracy. Modifications introduced at the phosphodiester bond (M4), such as with alpha-thio nucleoside triphosphates, have the advantage that these modifications
5 do not interfere with accurate Watson-Crick base-pairing and additionally allow for the one-step post-synthetic site-specific modification of the complete nucleic acid molecule e.g. via alkylation reactions [K.L. Nakamaye, G. Gish, F. Eckstein and H.-P. Vossberg, *Nucleic Acids Res.*, 16, 9947-59 (1988)]. However, this modification is not applicable where the
10 exonucleolytically released nucleotides are to be treated with immobilized alkaline phosphatase intermediate release and mass spectrometric detection. Mass modification can also occur at the sugar moiety, such as at the position C-2' (M3). Modifications at this position can be introduced with the purpose of modulating the rate of exonuclease activity in order to synchronize the degradation process from time to time. The modification M4 can also serve this purpose. For example, it is known [Burgers and Eckstein, (1979), *supra*] that a
15 phosphodiester bond carrying a monothio function is approximately 100 time less sensitive towards exonucleolytic degradation by exonuclease III.

The tables in FIGURES 7 and 8 depict some examples of mass-modifying functionalites for nucleotides. This list is, however, not meant to be limiting, since numerous other combinations of mass modifying functions and positions within the nucleotide
20 molecule are possible and are deemed part of the invention. The mass modifying functionality can be, for example, a halogen, an azido, or of the type XR. Wherein X is a linking group and R is a mass modifying functionality. The mass modifying functionality can thus be used to introduce defined mass increments into the nucleotide molecule.

Without limiting the scope of the invention, the mass modification, M, can be
25 introduced for X in XR as well as using oligo-/polyethylene glycol derivatives for R. The mass modifying increment in this case is 44, i.e., five different mass modified species could be generated by just changing m from 0 to 4 thus adding mass units of 45 (m=0), 89 (m=1), 133 (m=2), 177 (m=3) and 221 (m=4). The oligo/polyethylene glycols can also be monoalkylated by a lower alkyl such as methyl, ethyl, propyl, isopropyl, t-butyl and the like.
30 A selection of linking functionalities X are also illustrated in FIGURE 8. Other chemistries can be used in the mass modified compounds, as for example, those described recently in *Oligonucleotides and Analogues, A Practical Approach*, F. Eckstein, editor, IRL Press, Oxford, 1991.

In yet another embodiment, various mass modifying functionalities R, other
35 than oligo/polyethylene glycols, can be selected and attached via appropriate linking chemistries X. A simple mass modification can be achieved by substituting H for halogens like F, Cl, Br and/or I; or pseudohalogens such as SCN, NCS; or by using different alkyl and

$\text{Si}(\text{CH}_3)_2(\text{C}_2\text{H}_5)$, $\text{Si}(\text{CH}_3)(\text{C}_2\text{H}_5)_2$, $\text{Si}(\text{C}_2\text{H}_5)_3$. Yet another mass modification can be obtained by attaching homo- or heteropeptides through X to the nucleotide. One example useful in generating mass modified species with a mass increment of 57 is the attachment of oligoglycines, e.g. mass modifications of 74 ($r=1$, $m=0$), 131 ($r=1$, $m=2$), 188 ($r=1$, $m=3$), 245 ($r=1$, $m=4$) are achieved. Simple oligoamides also could be used, e.g. mass modifications of 74 ($r=1$, $m=0$), 88 ($r=2$, $m=0$), 102 ($r=3$, $m=0$), 116 ($r=4$, $m=0$) etc. are obtainable. For those skilled in the art it will be obvious that there are numerous possibilities, for introducing, in a predetermined manner, many different mass modifying functionalities to the nucleotide.

In yet another embodiment of this invention, the mass modifying functionality can be introduced by a two or multiple step process. In this case the nucleotide is, in a first step, modified by a precursor functionality such as azido, $-\text{N}_3$, or modified with a functional group in which the R in XR is H thus providing temporary functions e.g. but not limited to $-\text{OH}$, $-\text{NH}_2$, $-\text{NHR}$, $-\text{SH}$, $-\text{NCS}$, $-\text{OCO}(\text{CH}_2)_r\text{COOH}$ ($r = 1-20$), $-\text{NHCO}(\text{CH}_2)_r\text{COOH}$ ($r = 1-20$), $-\text{OSO}_2\text{OH}$, $-\text{OCO}(\text{CH}_2)_r\text{I}$ ($r = 1-20$), $-\text{OP}(\text{O-Alkyl})\text{N}(\text{Alkyl})_2$. These less bulky functionalities result in better substrate properties for enzymatic DNA or RNA synthesis reactions. The appropriate mass modifying functionality can then be introduced after the generation of the target nucleic acid prior to mass spectrometry and either prior to exonuclease degradation or after release by exonuclease action.

(iii) The exonuclease sequencer:

A schematic outlay of an exonuclease sequencer is shown in FIGURE 9. The central part is the reactor S which has a cooling/heating mantle (2) and a frit or semipermeable membrane (4). Several flasks (R1-R5) can be dedicated to supply reagents such as buffer solutions, enzyme, etc. through a cooling/heating coil T. Beneath the reactor S there is a capillary tube E in which either the exonuclease or the nucleic acids can be immobilized. It is within the scope of this invention that there are at least two different modes by which the system can be operated. In one mode, the nucleic acids are immobilized on beads or flat membrane disks placed in the reactor S, or alternatively, immobilized on the inner surface of the walls within the capillary E. Exonuclease is added to the reactor in a controlled manner and the reaction mixture circulated through a loop maintained at a carefully controlled temperature. In a second mode, the exonuclease can be immobilized in e.g. a capillary E beneath the reactor S or could be immobilized on beads or on a membrane or entrapped in a gel or kept in the reactor S by way of a semipermeable membrane. By varying the length and diameter of the capillary and the flow rate through a pumping device P, the contact time of the nucleic acids with the exonuclease can be varied.

immobilized alkaline phosphatase or other mass-modifying reagents. In case the liquid volume which is transferred to the mass spectrometer is too large, only a portion can be supplied while the remainder is separated from the flow stream by using a flow splitting device SP. Unused or excess solutions can be disposed of by the waste container W. In case
5 the reaction mixture of the exonuclease digestion is processed via a moving belt the liquid flow can be directed through this module prior to entering the mass spectrometer.

(iv) The exonuclease sequencing process:

10 Various exonucleases can be used, such as snake venom phosphodiesterase, Bal-31 nuclease, E. coli exonuclease VII, Mung Bean Nuclease, S1 Nuclease, exonuclease and exonuclease III as well as the exonuclease activity of some DNA polymerases such as E. coli DNA polymerase I, the Klenow fragment of DNA polymerase I, T4 or T7 DNA
15 polymerases, Taq DNA polymerase, Deep Vent DNA polymerase, and Vent_r DNA polymerase. The activity of these exonucleases can be modulated, for instance, by shifting off the optimal pH and/or temperature range or by adding poisoning agents to the reaction mixture. The exonuclease activity can also be modulated by way of functional groups, such as at the C-2' position of the sugar moiety of the nucleotide building block or at the
20 phosphodiester bond (i.e., M3/M4 in FIGURE 5 and 6).
In the instance that unmodified nucleotides are detected, the masses for the phosphate dianion are 329.209 for pdG, 313.210 for pdA, 304.196 for pdT and 289.185 for pdC. In an idealized system the enzymatic digestion would be initiated at all nucleic acid chains at the same time, and the nucleotides would be released in identical time intervals d and detected by their individual molecular weights one after the other by the mass
25 spectrometer. FIGURE 10 illustrates the signals versus time for the sequence 5'...A-T*-C-C-G-G-A 3'.

The influence of an activity modulating functionality M3/M4 on appropriately modified thymidine, T*, is also depicted. Due to the drastically reduced cleavage rate of the phosphodiester bond between dC and dT* the molecular mass representing the T* signal
30 appears after a longer time interval f. The significant retardation of the cleavage rate of one type of nucleotide results in better overall synchronization of the enzymatic process. This retardative effect can be varied to a large extent by the bulkiness of the modifying functional group as well as by a possible interaction with the active site of the exonuclease.
35 Additionally, partial overlap of signals can be resolved by computational methods.

(v) Multiplex exonuclease sequencing:

20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 251

FIGURE 11. For multiplex mass spectrometric exonuclease DNA sequencing, the DNA fragments to be processed in parallel can be identified by fragment specific labels introduced by the mass modifying functionality M. The target nucleic acid sequences can be mass-modified by using, for example, unmodified dNTP⁰s or NTP⁰s (Tr⁰), nucleoside triphosphates mass-modified with the same functional group, such as an additional methyl group at the heterocyclic base, using either dNTP¹ or NTP¹ (Tr¹), mass difference large enough to be discriminated from the nucleotides of Tr⁰ or Tr¹, such as e.g. an ethyl group at the heterocyclic base, by employing either dNTP² or NTP² etc. Thus i modified DNA fragments can be simultaneously exonuclease sequenced. For example, the i different DNA fragments can be immobilized on different membranes and a stack of such membranes placed into the reactor S (FIGURE 9) for simultaneous exonuclease mass spectrometric sequencing. Since the individual molecular weights of the various mass-modified four nucleotides are known in advance, the mass spectrometrically detected nucleotide masses can be easily assigned to the parent nucleic acid fragments and thus several sequences can be detected simultaneously by the mass spectrometer. Even in the worst case when the same nucleotide, e.g. dT is at the same position in all sequences, processed in parallel the signal can be decoded due to the difference in mass between dT⁰, dT¹, dT², dT³, ..., dTⁱ.

The synchronization of exonuclease action can be improved by incorporating modified nucleotides (modified at C-2' or at the phosphodiester bond) into the otherwise unmodified or mass-modified nucleic acid fragments as set out above. In particular such a mass-modified nucleotide can also be introduced at the 3' end of the single stranded nucleic acid fragment (i.e., C-2' or phosphodiester bond modifications) to achieve a more uniform initiation of the exonuclease reaction.

In yet another embodiment of this invention, a reduction of overlap of neighboring signals can be achieved by using a moving belt device as schematically shown in FIGURE 12. In a recent publication a moving belt has been described although in a completely unrelated application [M.Moini and F.P. Abramson, Biological Mass Spectrometry, 20, 308-12 (1991)]. The effect of the moving belt can be to help spread the appearance of sequentially released nucleotide/nucleoside signals as illustrated in FIGURE 13. The width between consecutive, i.e., neighboring, signals can be increased with the speed of the moving belt.

Without limiting the scope of the invention FIGURE 12 illustrates a possible configuration of the moving belt module for exonuclease mediated mass spectrometric sequencing. An endless metal ribbon (10) is driven with variable speed using a controllable stepping motor and appropriately positioned pulleys. Spring-loaded pulleys can be used to maintain a sufficiently high tension on the moving belt. The sample is applied to the belt from the reactor module A (FIGURE 9) at a position which is in direct contact with a cooling/heating plate B.

onto the belt. Crystal formation can be observed with a viewing device D (CCD camera and optics). Alternatively the container M can be used to mix the sample with a diluent or other reagent, enzyme, internal standard etc. at a mixing valve or vortex (12). In the instance of relatively small molecules such as the released nucleotides, matrix is not essential for the laser desorption/ionization process to take place. The belt 10 moves the sample under a laser source E (with appropriate optics) for desorption and ionization.

A heating element C, such as a microwave source, can be placed near the surface of the returning belt, separated from the forward moving belt by an insulating shield I, to clean the surface of the metal ribbon belt 10 of any organic material prior to loading a new sample. Alternatively, a washing station W can be integrated before the heating element C; in this case the function of the heating element C can be completely dry the metal ribbon prior to reloading of sample. Before and after the laser targets the sample, two differential vacuum pumping stages F and G are positioned. An electric field is applied after the second vacuum stage to accelerate the ions into the mass spectrometer. As mass analyzer, a quadrupole can be used, though other mass analyzing configurations are known in the art and are within the scope of the invention. The design of the vacuum interface of the moving belt between the sample application compartment which is at atmospheric pressure, and the mass spectrometer can be important. In one approach, this vacuum seal can be provided by the use of tunnel seals in a two-stage vacuum lock as previously described [Moini et al, (1991), *supra*].

As described above, an increase of throughput can be obtained by multiplexing. In yet another embodiment of the invention the moving belt device can be used for a second dimension in multiplexing by applying s samples from s sequencing reactors A (FIGURE 9) simultaneously in different locations onto the moving belt. Desorption and ionization of these multiple samples is achieved by moving the laser beam with adjustable speed and frequency back and forth perpendicular to the direction of the moving belt. Identification and assignment of the nucleotides/nucleosides detected to the various nucleic acid fragments can be achieved by second dimension multiplexing, i.e., by mass-labeling the nucleic acid fragments with for example $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$ and $-\text{CH}_2(\text{CH}_2)_r\text{CH}_2-$, and labeling the different reactors, for example, by individual halogen atoms, i.e., F for reactor 1 (thus having the different DNA fragments labeled with $-\text{CH}_2\text{F}$, $-\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2(\text{CH}_2)_r\text{CH}_2\text{F}$), Cl for reactor 2 (thus having the different DNA fragments labeled with $-\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2(\text{CH}_2)_r\text{CH}_2\text{Cl}$), Br for reactor 3 etc. This can increase the throughput dramatically. Two-dimensional multiplexing can be applied in different ways. For example, it can be used to simultaneously sequence the fragments forming one set of overlapping deletions of the same long DNA insert. In another embodiment, several subsets of ordered deletions of different DNA inserts

The enormous advantage of exonuclease mediated mass spectrometric DNA sequencing is that small molecules are analyzed and identified by mass spectrometry. In this mass range, the accuracy of mass spectrometers is routinely very high, i.e., 0.1 mass units are easily detected. This increases the potential for multiplexing as small differences in mass can be detected and resolved. An additional advantage of mass spectrometric sequencing is that the identified masses can be registered automatically by a computer and by adding the time coordinate automatically aligned to sequences. Since the sequences so determined are memorized (i.e., saved to disk or resident in the computer memory) appropriate existing computer programs operating in a multitasking environment can be searching in the "background" (i.e., during continuous generation of new sequence data by the exonuclease mass spectrometric sequencer) for overlaps and generate contiguous sequence information which, via a link to a sequence data bank, can be used in homology searches, etc.

Another aspect of this invention concerns kits for sequencing nucleic acids by exonuclease mass spectrometry, which include combinations of the above described sequencing reactants. For instance, in one embodiment, the kit comprises reagents for multiplex mass spectrometric sequencing of several different species of nucleic acid. The kit can include an exonuclease for cleaving the nucleic acids unilaterally from a first end to sequentially release individual nucleotides, a set of nucleotides for synthesizing the different species of nucleic acids, at least a portion of the nucleotides being mass-modified such that sequentially released nucleotides of each of the different species of nucleic acids are distinguishable, a polymerase for synthesizing the nucleic acids from complementary templates and the set of nucleotides, and a solid support for immobilizing one of the nucleic acids or the exonuclease. The kit can also include appropriate buffers, as well as instructions for performing multiplex mass spectrometry to concurrently sequence multiple species of nucleic acids. In another embodiment, the sequencing kit can include an exonuclease for cleaving a target nucleic acid unilaterally from a first end to sequentially release individual nucleotides, a set of nucleotides for synthesizing the different species of nucleic acids, at least a portion of the nucleotides being mass-modified to modulate the exonuclease activity, a polymerase for synthesizing the nucleic acid from a complementary template and the set of nucleotides, and a solid support for immobilizing one of the nucleic acids or the exonuclease.

Another aspect of this invention concerns a "reverse-Sanger" type sequencing method using exonuclease digestion of nucleic acids to produce a ladder of nested digestion fragments detectable by mass spectrometry. For instance, as above, the target nucleic acid can be immobilized to a solid support to provide unilateral degradation of the chain by exonuclease action. Incorporating into the target nucleic acid a limited number of mass-modified nucleotides which inhibit the exonuclease activity (i.e., protect an individual nucleic acid chain from further degradation) can result in a ladder of nested exonuclease fragments

from the solid support (i.e., via a cleavable linkage) and the molecular weight values for each species of the nested fragments determined by mass spectrometry, as described in U.S. Patent application No. 08/001,323. From the molecular weight values determined, the sequence of the nucleic acid can be generated. It is clear that many variations of this reaction are possible and that it is amenable to multiplexing. For example, the target nucleic acid need not be bound to a solid support, rather any protecting group can be used to ensure unilateral exonuclease degradation. Where mass-modified nucleotides are used which have large enough molecular weight differences to be differentiated between by mass spectrometry (i.e., the termination of a chain with a particular mass-modified nucleotide is discernable from all other terminations), the exonuclease sequencing can be carried out to create only one set of nested fragments. Alternatively, individual types of exonuclease-inhibiting nucleotides can be incorporated in separate reactions to create sets of nested fragments. For instance, four sets of nested fragments can be separately generated wherein one set terminates with mass-modified A's, one set terminates in mass-modified G's, etc. and the total sequence is determined by aligning the collection of nested exonuclease fragments.

EXAMPLE 1

Immobilization of nucleic acids to solid supports via disulfide bonds.

As a solid support, SEQUELON membranes (Millipore Corp., Bedford, MA) with phenyl isothiocyanate groups is used as a starting material. The membrane disks, with a diameter of 8 mm, are wetted with a solution of N-methylmorpholine/water/2-propanol (NMM solution) (2/49/49;v/v/v), the excess liquid removed with filterpaper and placed on a piece of plastic film or aluminium foil located on a heating block set to 55°C. A solution of 1 mM 2-mercaptoethylamine (cysteamine) or 2,2'-dithio-bis(ethylamine) (cystamine) or S-(2-thiopyridyl)-2-thio-ethylamine (10 ul, 10 nmol) in NMM is added per disk and heated at 55°C. After 15 min 10 ul of NMM solution are added per disk and heated for another 5 min. Excess of isothiocyanate groups may be removed by treatment with 10 ul of a 10 mM solution of glycine in NMM solution. In case of cystamine the disks are treated with 10 ul of a solution of 1M aqueous dithiothreitol (DTT)/2-propanol (1:1, v/v) for 15 min at room temperature. Then the disks are thoroughly washed in a filtration manifold with 5 aliquots of 1 ml each of the NMM solution, then with 5 aliquots of 1 ml acetonitrile/water (1/1; v/v) and subsequently dried. If not used immediately the disks are stored with free thiol groups in a solution of 1M aqueous dithiothreitol/2-propanol (1:1; v/v) and, before use, DTT is removed by three washings with 1ml each of the NMM solution. Single-stranded nucleic acid fragments with 5'-SH functionality can be prepared by various methods [e.g. B.C.F Chu et al., Nucl eic Acids Res., 14, 5591-5603 (1986), Sproat et al., Nucleic Acids Res., 15, 4837-48 (1987) and Oligonucleotides and Analogues. A Practical Approach (F. Eckstein editor) IRL

... under oxidizing conditions in general.

it is sufficient to add the 5'-thiolated nucleic acid fragments dissolved in 10 ul 10 mM de-aerated triethylammonium acetate buffer (TEAA) pH 7.2 to the thiolated membrane supports: coupling is achieved by drying the samples onto the membrane disks with a cold fan. This process can be repeated by wetting the membrane with 10 ul of 10 mM TEAA buffer pH 7.2 and drying as before. When using the 2-thiopyridyl derivatized compounds anchoring can be monitored by the release of pyridine-2-thione spectrophotometrically at 343 nm.

In another variation of this approach the single-stranded nucleic acid is functionalized with an amino group at the 5'-end by standard procedures. The primary amino group is reacted with 3-(2-pyridyldithio)propionic acid N-hydroxysuccinimide ester (SPDP) and subsequently coupled to the thiolated supports and monitored by the release of pyridyl-2-thione as described above. After denaturation of any remaining protein and ethanol precipitation of the functionalized nucleic acid, the pellet is dissolved in 10 ul 10 mM TEAA buffer pH 7.2 and 10 ul of a 2 mM solution of SPDP in 10 mM TEAA are added. The reaction mixture is vortexed and incubated for 30 min at 25°C; excess SPDP is then removed by three extractions (vortexing, centrifugation) with 50 ul each of ethanol and the resulting pellets dissolved in 10 ul 10 mM TEAA buffer pH 7.2 and coupled to the thiolated supports (see above).

The immobilized nucleic acids can be released by three successive treatments with 10 ul each of 10 mM 2-mercaptoethanol in 10 mM TEAA buffer pH 7.2.

EXAMPLE 2

Immobilization of nucleic acids on solid support via a levulinyl group.

5-Aminolevulinic acid is protected at the primary amino group with the Fmoc group using 9-fluorenylmethyl N-succinimidyl carbonate and then transformed into the N-hydroxysuccinimide ester (NHS ester) using N-hydroxysuccinimide and dicyclohexyl carbodiimide under standard conditions. Nucleic acids which are functionalized with primary amino acid at the 5' end are EtOH precipitated and resuspended in 10 ul of 10 mM TEAA buffer pH 7.2. 10 ul of a 2 mM solution of the Fmoc-5-aminolevulinyl-NHS ester in 10 mM TEAA buffer is added, vortexed and incubated at 25°C for 30 min. The excess of reagents can be removed by ethanol precipitation and centrifugation. The Fmoc group is cleaved off by resuspending the pellets in 10 ul of a solution of 20% piperidine in N,N-dimethylformamide/water (1:1, v/v). After 15 min at 25°C piperidine is thoroughly removed by three precipitations/centrifugations with 100 ul each of ethanol, the pellets resuspended in 10 ul of a solution of N-methylmorpholine, propanol-2 and water (2/10/88; v/v/v) and coupled to the solid support carrying an isothiocyanate group. In case of the DITC-Sequelon membrane (Millipore Corp., Bedford, MA) the membranes are prepared as described in EXAMPLE 1 and coupling is achieved on a heating block at 55°C as described above. The procedure can be applied to...

The immobilized nucleic acids can be released from the solid support by three successive treatments with 10 ul of 100 mM hydrazinium acetate buffer pH 6.5.

EXAMPLE 3

5 Immobilization of nucleic acids on solid supports via a trypsin sensitive linkage.

Sequelon DITC membrane disks of 8 mm diameter (Millipore Corp. Bedford, MA) are wetted with 10 ul of NMM solution (N-methylmorpholine/propanol-2/water; 2/49/49; v/v/v) and a linker arm introduced by reaction with 10 ul of a 10 mM solution of 1,6-diaminohexane in NMM. The excess of the diamine is removed by three washing steps with 100 ul of NMM solution. Using standard peptide synthesis protocols two L-lysine residues are attached by two successive condensations with N-Fmoc-N-tBoc-L-lysine pentafluorophenylester, the terminal Fmoc group is removed with piperidine in NMM and the free e-amino group coupled to 1,4-phenylene diisothiocyanate (DITC). Excess DITC is removed by three washing steps with 100 ul propanol-2 each and the N-tBoc groups removed with trifluoroacetic acid according to standard peptide synthesis procedures. The nucleic acids are prepared from as above from a primary amino group at the 5'-terminus. The ethanol precipitated pellets are resuspended in 10 ul of a solution of N-methylmorpholine, propanol-2 and water (2/10/88; v/v/v) and transferred to the Lys-Lys-DITC membrane disks and coupled on a heating block set at 55°C. After drying 10 ul of NMM solution is added and the drying process repeated.

The immobilized nucleic acids can be cleaved from the solid support by treatment with trypsin.

EXAMPLE 4

25 Immobilization of nucleic acids on solid supports via pyrophosphate linkage.

The DITC Sequelon membrane (disks of 8 mm diameter) are prepared as described in EXAMPLE 3 and 10 ul of a 10 mM solution of 3-aminopyridine adenine dinucleotide (APAD) (Sigma) in NMM solution added. The excess of APAD is removed by a 10 ul wash of NMM solution and the disks are treated with 10 ul of 10 mM sodium periodate in NMM solution (15 min, 25°C). Excess of periodate is removed and the having a primary amino group at the 5'-end are dissolved in 10 ul of a solution of N-methylmorpholine/propanol-2/water (2/10/88; v/v/v) and coupled to the 2',3'-dialdehyde functions of the immobilized NAD analog.

The immobilized nucleic acids can be released from the solid support by treatment with either NADase or pyrophosphatase in 10 mM TEAA buffer at pH 7.2 at 37°C for 15 min.

EXAMPLE 5

Synthesis of pyrimidine nucleotides mass modified at C-5 of the heterocyclic base with glycine residues.

Starting material is 5-(3-aminopropynyl-1)-3',5'-di-p-tolyldeoxyuridine prepared and 3',5'-de-O-acylated according to literature procedures [Haralambidis et al., Nucleic Acids Res., 15, 4857-76 (1987)]. 0.281 g (1.0 mmole) 5-(3-aminopropynyl-1)-2'-deoxyuridine are reacted with 0.927 g (2.0 mmole) N-Fmoc-glycine pentafluorophenylester in 5 ml absolute N,N-dimethylformamide in the presence of 0.129 g (1 mmole; 174 μ l) N,N-diisopropylethylamine for 60 min at room temperature. Solvents are removed by rotary evaporation and the product purified by silica gel chromatography (Kieselgel 60, Merck; column: 2.5x 50 cm, elution with chloroform/methanol mixtures). Yield 0.44 g (0.78 mmole, 78 %). In order to add another glycine residue the Fmoc group is removed with a 20 min treatment with 20 % solution of piperidine in DMF, evaporated in vacuo and the remaining solid material extracted three times with 20 ml ethylacetate; after having removed the remaining ethylacetate N-Fmoc-glycine pentafluorophenylester is being coupled as described above. This glycine modified thymidine analogue building block for chemical DNA synthesis can be used to substitute for thymidine or uridine nucleotides in the target nucleic acid.

EXAMPLE 6

Synthesis of pyrimidine nucleotides mass modified at C-5 of the heterocyclic base with β -alanine residues.

Starting material is the same as in EXAMPLE 5. 0.281 g (1.0 mmole) 5-(3-Aminopropynyl-1)-2'-deoxyuridine is reacted with N-Fmoc- β -alanine pentafluorophenylester (0.955 g, 2.0 mmole) in 5 ml N,N-dimethylformamide (DMF) in the presence of 0.129 g (174 μ l; 1.0 mmole) N,N-diisopropylethylamine for 60 min at room temperature. Solvents are removed and the product purified by silica gel chromatography as described in EXAMPLE 6. Yield: 0.425 g (0.74 mmole, 74 %). Another β -alanine moiety could be added in exactly the same way after removal of the Fmoc group. This building block can be substitute for any of the thymidine or uridine residues in the target nucleic acid.

EXAMPLE 7

Synthesis of a pyrimidine nucleotide mass modified at C-5 of the heterocyclic base with ethylene glycol monomethyl ether.

As nucleosidic component 5-(3-aminopropynyl-1)-2'-deoxyuridine is used in this example (see EXAMPLE 5 and 6). The mass-modifying functionality is obtained as follows: 7.61 g (100.0 mmole) freshly distilled ethylene glycol monomethyl ether dissolved in 50 ml absolute N,N-dimethylformamide (DMF) is reacted with 0.129 g (1 mmole; 174 μ l) N,N-diisopropylethylamine for 60 min at room temperature. Solvents are removed by rotary evaporation and the product purified by silica gel chromatography (Kieselgel 60, Merck; column: 2.5x 50 cm, elution with chloroform/methanol mixtures). Yield 0.44 g (0.78 mmole, 78 %).

room temperature. The reaction is terminated by the addition of water (5.0 ml), the reaction mixture evaporated in vacuo, co-evaporated twice with dry toluene (20 ml each) and the residue redissolved in 100 ml dichloromethane. The solution is extracted successively, twice with 10 % aqueous citric acid (2 x 20 ml) and once with water (20 ml) and the organic phase dried over anhydrous sodium sulfate. The organic phase is evaporated in vacuo, the residue redissolved in 50 ml dichloromethane and precipitated into 500 ml pentane and the precipitate dried in vacuo. Yield: 13.12 g (74.0 mmole; 74 %). 8.86 g (50.0 mmole) of succinylated ethylene glycol monomethyl ether is dissolved in 100 ml dioxane containing 5 % dry pyridine (5 ml) and 6.96 g (50.0 mmole) 4-nitrophenol and 10.32 g (50.0 mmole) dicyclohexylcarbodiimide is added and the reaction run at room temperature for 4 hours. Dicyclohexylurea is removed by filtration, the filtrate evaporated in vacuo and the residue redissolved in 50 ml anhydrous DMF. 12.5 ml (about 12.5 mmole 4-nitrophenylester) of this solution is used to dissolve 2.81 g (10.0 mmole) 5-(3-aminopropynyl-1)-2'-deoxyuridine. The reaction is performed in the presence of 1.01 g (10.0 mmole; 1.4 ml) triethylamine at room temperature overnight. The reaction mixture is evaporated in vacuo, co-evaporated with toluene, redissolved in dichloromethane and chromatographed on silicagel (Si60, Merck; column 4x50 cm) with dichloromethane/methanol mixtures. The fractions containing the desired compound are collected, evaporated, redissolved in 25 ml dichloromethane and precipitated into 250 ml pentane.

EXAMPLE 8

Synthesis of pyrimidine nucleotides mass-modified at C-5 of the heterocyclic base with diethylene glycol monomethyl ether.

Nucleosidic starting material is as in previous examples, 5-(3-aminopropynyl-1)-2'-deoxyuridine. The mass-modifying functionality is obtained similar to EXAMPLE 7. 12.02 g (100.0 mmole) freshly distilled diethylene glycol monomethyl ether dissolved in 50 ml absolute pyridine is reacted with 10.01 g (100.0 mmole) recrystallized succinic anhydride in the presence of 1.22 g (10.0 mmole) 4-N,N-dimethylaminopyridine (DMAP) overnight at room temperature. The work-up is as described in EXAMPLE 7. Yield: 18.35 g (82.3 mmole, 82.3 %). 11.06 g (50.0 mmole) of succinylated diethylene glycol monomethyl ether is transformed into the 4-nitrophenylester and subsequently 12.5 mmole reacted with 2.81 g (10.0 mmole) of 5-(3-aminopropynyl-1)-2'-deoxyuridine as described in EXAMPLE 7. Yield after silica gel column chromatography and precipitation into pentane: 3.34 g (6.9 mmole, 69 %).

EXAMPLE 9

Synthesis of deoxyadenosine mass-modified at C-8 of the heterocyclic base with glycine.

Starting material is N6-benzoyl-8-bromo-5'-O-(4,4'-dimethoxytrityl)-2'-deoxyadenosine prepared according to literature [Singh et al., Nucleic Acids Res. 18, 3339-45 (1990)]. 632.5 mg (1.0 mmole) of this 8-bromo-deoxyadenosine derivative is suspended in 5 ml absolute ethanol and reacted with 251.2 mg (2.0 mmole) glycine methyl ester (hydrochloride) in the presence of 241.4 mg (2.1 mmole; 366 ul) N,N-diisopropylethylamine and refluxed until the starting nucleosidic material has disappeared (4-6 hours) as checked by thin layer chromatography (TLC). The solvent is evaporated and the residue purified by silica gel chromatography (column 2.5x50 cm) using solvent mixtures of chloroform/methanol containing 0.1 % pyridine. The product fractions are combined, the solvent evaporated, dissolved in 5 ml dichloromethane and precipitated into 100 ml pentane. Yield: 487 mg (0.76 mmole, 76 %).

EXAMPLE 10

Synthesis of deoxyadenosine mass-modified at C-8 of the heterocyclic base with glycylglycine.

This derivative is prepared in analogy to the glycine derivative of EXAMPLE 9. 632.5 mg (1.0 mmole) N6-Benzoyl-8-bromo-5'-O-(4,4'-dimethoxytrityl)-2'-deoxyadenosine is suspended in 5 ml absolute ethanol and reacted with 324.3 mg (2.0 mmole) glycyl-glycine methyl ester in the presence of 241.4 mg (2.1 mmole, 366 ul) N,N-diisopropylethylamine. The mixture is refluxed and completeness of the reaction checked by TLC. Work-up and purification is similar as described in EXAMPLE 9. Yield after silica gel column chromatography and precipitation into pentane: 464 mg (0.65 mmole, 65 %).

EXAMPLE 11

Synthesis of deoxythymidine mass-modified at the C-2' of the sugar moiety with ethylene glycol monomethyl ether residues.

Starting material is 5'-O-(4,4-dimethoxytrityl)-2'-amino-2'-deoxythymidine synthesized according to published procedures [e.g. Verheyden et al., J. Org. Chem., 36, 250-254 (1971); Sasaki et al., J. Org. Chem., 41, 3138-3143 (1976); Imazawa et al., J. Org. Chem., 44, 2039-2041 (1979); Hobbs et al., J. Org. Chem., 42, 714-719 (1976); Ikehara et al., Chem. Pharm. Bull. Japan, 26, 240-244 (1978); see also PCT Application WO 88/00201]. 5'-O-(4,4-Dimethoxytrityl)-2'-amino-2'-deoxythymidine (559.62 mg; 1.0 mmole) is reacted with 2.0 mmole of the 4-nitrophenyl ester of succinylated ethylene glycol monomethyl ether (see EXAMPLE 7) in 10 ml dry DMF in the presence of 1.0 mmole (140 ul) triethylamine for 18 hours at room temperature. The reaction mixture is evaporated in vacuo, co-evaporated with

triethylamine). The product containing fractions are combined, evaporated and precipitated into pentane. Yield: 524 mg (0.73 mmol; 73 %).

In an analogous way, employing the 4-nitrophenyl ester of succinylated diethylene glycol monomethyl ether (see EXAMPLE 8) and triethylene glycol monomethyl ether the corresponding mass modified deoxythymidine is prepared. The mass difference between the ethylene, diethylene and triethylene glycol derivatives is 44.05, 88.1 and 132.15 dalton respectively.

EXAMPLE 12

Synthesis of deoxyuridine-5'-triphosphate mass-modified at C-5 of the heterocyclic base with glycine, glycyl-glycine and β -alanine residues.

0.281 g (1.0 mmole) 5-(3-Aminopropynyl-1)-2'-deoxyuridine (see EXAMPLE 5) is reacted with either 0.927 g (2.0 mmole) N-Fmoc-glycine pentafluorophenylester or 0.955g (2.0 mmole) N-Fmoc- β -alanine pentafluorophenyl ester in 5 ml dry DMF in the presence of 0.129 g N,N-diisopropylethylamine (174 μ l, 1.0 mmole) overnight at room temperature. Solvents are removed by evaporation in vacuo and the condensation products purified by flash chromatography on silica gel [Still et al., J. Org. Chem., 43, 2923-2925 (1978)]. Yields: 476 mg (0.85 mmole; 85 %) for the glycine and 436 mg (0.76 mmole; 76 %) for the β -alanine derivative. For the synthesis of the glycyl-glycine derivative the Fmoc group of 1.0 mmole Fmoc-glycine-deoxyuridine derivative is removed by one-hour treatment with 20 % piperidine in DMF at room temperature. Solvents are removed by evaporation in vacuo, the residue is co-evaporated twice with toluene and condensed with 0.927 g (2.0 mmole) N-Fmoc-glycine pentafluorophenyl ester and purified following standard protocol. Yield: 445 mg (0.72 mmole; 72 %). The glycyl-, glycyl-glycyl- and β -alanyl-2'-deoxyuridine derivatives N-protected with the Fmoc group are now transformed to the 3'-O-acetyl derivatives by tritylation with 4,4-dimethoxytrityl chloride in pyridine and acetylation with acetic anhydride in pyridine in a one-pot reaction and subsequently detritylated by one-hour treatment with 80 % aqueous acetic acid according to standard procedures. Solvents are removed, the residues dissolved in 100 ml chloroform and extracted twice with 50 ml 10 % sodium bicarbonate and once with 50 ml water, dried with sodium sulfate, the solvent evaporated and the residues purified by flash chromatography on silica gel. Yields: 361 mg (0.60 mmole; 71 %) for the glycyl-, 351 mg (0.57 mmole; 75 %) for the β -alanyl- and 323 mg (0.49 mmole; 68 %) for the glycyl-glycyl-3'-O'-acetyl-2'-deoxyuridine derivatives respectively. Phosphorylation at the 5'-OH with POCl₃, transformation into the 5'-triphosphate by in-situ reaction with tetra(tri-n-butylammonium) pyrophosphate in DMF, 3'-de-O-acetylation and cleavage of the Fmoc group and final purification by anion-exchange chromatography on DEAE-Sephadex is performed. Yields according to UV-absorbance of

mmole (75 %) and 5-(3-(N-glycyl-glycyl)-amidopropynyl-1)-2'-deoxyuridine- 5'-triphosphate 0.38 mmole (78 %).

EXAMPLE 13

5 Synthesis of 8-glycyl- and 8-glycyl-glycyl-2'-deoxyadenosine-5'-triphosphate.

727 mg (1.0 mmole) of N6-(4-tert.butylphenoxyacetyl)-8-glycyl-5'-(4,4-dimethoxytrityl)-2'- deoxyadenosine or 800 mg (1.0 mmole) N6-(4-tert.butylphenoxyacetyl)-8-glycyl-glycyl-5'-(4,4-dimethoxytrityl)-2'-deoxyadenosine prepared according to EXAMPLES 9 and 10 and literature [Koster et al., Tetrahedron, 37 : 362 (1981)] are
10 acetylated with acetic anhydride in pyridine at the 3'-OH, detritylated at the 5'-position with 80 % acetic acid in a one-pot reaction and transformed into the 5'-triphosphates via phosphorylation with POCl₃ and reaction in-situ with tetra(tri-n-butylammonium) pyrophosphate. Deprotection of the N6-tert-butylphenoxyacetyl, the 3'-O-acetyl and the O-methyl group at the glycine residues is achieved with concentrated aqueous ammonia for
15 three hours at room temperature. Ammonia is removed by lyophilization and the residue washed with dichloromethane, solvent removed by evaporation in vacuo and the remaining solid material purified by anion-exchange chromatography on DEAE-Sephadex using a linear gradient of triethylammonium bicarbonate from 0.1 to 1.0 M. The nucleoside triphosphate containing fractions (checked by TLC on polyethyleneimine cellulose plates) are combined
20 and lyophilized. Yield of the 8-glycyl-2'-deoxyadenosine-5'-triphosphate (determined by the UV-absorbance of the adenine moiety) is 57 % (0.57 mmole); the yield for the 8-glycyl-glycyl-2'-deoxyadenosine-5'-triphosphate is 51 % (0.51 mmole).

25 All of the above-cited references and publications are hereby incorporated by reference.

Equivalents

30 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the following claims.

- 29 -

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- 5
- (i) APPLICANT:
- (A) NAME: KOSTER, HUBERT
- (B) STREET: 1640 MONUMENT STREET
- (C) CITY: CONCORD
- 10 (D) STATE: MASSACHUSETTS
- (E) COUNTRY: USA
- (F) POSTAL CODE (ZIP): 01742
- (G) TELEPHONE: (508) 369-9790
- 15 (ii) TITLE OF INVENTION: DNA SEQUENCING BY MASS SPECTROMETRY
- (iii) NUMBER OF SEQUENCES: 6
- (v) COMPUTER READABLE FORM:
- 20 (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: ASCII text
- 25 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/034,738
- (B) FILING DATE: 19 March 1993
- (C) CLASSIFICATION:
- 30 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/034,738
- (B) FILING DATE: 19-MAR-1993
- (C) CLASSIFICATION:
- 35 (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: DeConti, Giulio A.
- (B) REGISTRATION NUMBER: 31,503
- (C) REFERENCE/DOCKET NUMBER: HKI-005PC
- 40 (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: (617) 227-7400
- (B) TELEFAX: (617) 227-5941
- 45 (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- 50 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- 55 (iii) HYPOTHETICAL: YES

- 30 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCCTTAGCTA

10

5 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

15 (iii) HYPOTHETICAL: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

20

GCGGCCGCAG GTCA

14

(2) INFORMATION FOR SEQ ID NO:3:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(iii) HYPOTHETICAL: YES

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AATTCAGCGG CCGC

14

40

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

45 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

50

(iii) HYPOTHETICAL: YES

- 31 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGCCGCAGGT CA

12

5

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

15

(iii) HYPOTHETICAL: YES

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGCCGCTGAA TT

12

25

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: other nucleic acid

35

(iii) HYPOTHETICAL: YES

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCTAACTTGC

10

Claims

1. A method of determining a sequence of a nucleic acid, comprising
 - (i) isolating the nucleic acid to be sequenced;
 - (ii) cleaving the nucleic acid unilaterally from a first end with an
5 exonuclease activity to sequentially release individual nucleotides;
 - (iii) identifying each of the sequentially release nucleotides by mass
spectrometry; and
 - (iv) determining the sequence of the nucleic acid from the identified
nucleotides.
- 10 2. The method of according to claim 1, wherein the nucleic acid is a 2'-
deoxyribonucleic acid (DNA).
3. The method of according to claim 1, wherein the nucleic acid is a ribonucleic
15 acid (RNA).
4. The method of according to claim 1, wherein the exonuclease is selected from
a group consisting of snake venom phosphodiesterase, spleen phosphodiesterase, Bal-31
nuclease, E. coli exonuclease I, E. coli exonuclease VII, Mung Bean Nuclease, S1 Nuclease,
20 an exonuclease activity of E. coli DNA polymerase I, an exonuclease activity of a Klenow
fragment of DNA polymerase I, an exonuclease activity of T4 DNA polymerase, an
exonuclease activity of T7 DNA polymerase, an exonuclease activity of Taq DNA
polymerase, an exonuclease activity of Deep Vent DNA polymerase, and an exonuclease
activity of Vent_r DNA polymerase.
- 25 5. The method according to claim 1, wherein the exonuclease is immobilized by
covalent attachment to a solid support, entrapment within a gel matrix, or contained in a
reactor with a semipermeable membrane.
- 30 6. The method according to claim 5, wherein the solid support is a capillary and
the exonuclease activity is covalently attached to an inner wall of the capillary.
7. The method according to claim 5, wherein the solid support is selected from a
group consisting of glass beads, cellulose beads, polystyrene beads, Sephadex beads,
35 Sepharose beads, polyacrylamide beads and agarose beads.
8. The method according to claim 5, wherein the solid support is a flat

9. The method according to claim 1, wherein the nucleic acid is immobilized by covalent attachment to a solid support and the exonuclease is in a solution contacted with the immobilized polynucleotide.
- 5 10. The method according to claim 9, wherein the solid support is a capillary and the nucleic acid is covalently attached to an inner wall of the capillary.
11. The method according to claim 9, wherein the solid support is selected from a group consisting of glass beads, cellulose beads, polystyrene beads, Sephadex beads,
10 Sepharose beads, polyacrylamide beads and agarose beads.
12. The method according to claim 9, wherein the solid support is a flat membrane.
- 15 13. The method according to claim 1, wherein the nucleic acid comprises mass-modified nucleotides.
14. The method according to claim 13, wherein the mass-modified nucleotides modulate the rate of the exonuclease activity.
20
15. The method according to claim 1, wherein the sequentially released nucleotides are mass-modified subsequent to exonuclease release and prior to mass spectrometric identification.
- 25 16. The method according to claim 15, wherein the sequentially released nucleotides are mass-modified by contact with an alkaline phosphatase.
17. The method according to claim 1, wherein i different species of nucleic acids are concurrently sequenced by multiplex mass spectrometric sequencing and the sequentially
30 released nucleotides from each species of the i nucleic acids can be distinguished by mass spectrometry from the sequentially released nucleotides from the remaining $i-1$ nucleic acids based on a difference in mass due to mass-modification of at least a portion of the sequentially released nucleotides.
- 35 18. The method according to claim 17, wherein the mass-modified nucleotides further serve to modulate activity of the exonuclease.

19. The method according to claim 17, wherein the mass-modified nucleotide comprises a mass-modifying functionality (M) attached to the sequentially released nucleotide.
- 5 20. The method according to claim 20, wherein at least one of the mass-modified nucleotides is modified with a mass modifying functionality (M) attached to a 5' phosphate.
21. The method according to claim 20, wherein at least one of the mass-modified nucleotides is modified with a mass modifying functionality (M) attached to a C-2' position
10 of a sugar moiety.
22. The method according to claim 20, wherein at least one of the mass-modified nucleotides is modified with a mass modifying functionality (M) attached to a heterocyclic base.
- 15 23. The method according to claim 22, wherein the mass modified nucleotide comprises a modified heterocyclic base selected from the group consisting of a cytosine moiety modified at C-5, an uracil moiety modified at C-5, a thymine moiety modified at the C-5 methyl group, an adenine moiety modified at C-8, a c7-deazaadenine moiety modified at C-8, a c7-deazaadenine moiety modified at C-7, a guanine moiety modified at C-8, a c7-deazaguanine moiety modified at C-8, a c7-deazaguanine moiety modified at C-7, a hypoxanthine moiety modified at C-8, a c7-deazahypoxanthine moiety modified at C-8 and a c7-deazahypoxanthine moiety modified at C-7.
- 20 24. The method according to claim 20, wherein the mass modifying functionality (M) is selected from a group consisting of XR, F, Cl, Br, I, Si(CH₃)₃, Si(CH₃)₂(C₂H₅), Si(CH₃)(C₂H₅)₂, Si(C₂H₅)₃, CH₂F, CHF₂, and CF₃, wherein X is selected from a group consisting of -OH, -NH₂, -NHR, -SH, -NCS, -OCO(CH₂)_rCOOH (where r = 1-20), -NHCO(CH₂)_rCOOH (where r = 1-20), -OSO₂OH, -OCO(CH₂)_rI (where r = 1-20), and -
25 OP(O-Alkyl)N(Alkyl)₂, and R is selected from a group consisting of H, methyl, ethyl, propyl, isopropyl, t-butyl, hexyl, benzyl, benzhydryl, trityl, substituted trityl, aryl, substituted aryl, polyoxymethylene, monoalkylated polyoxymethylene, a polyethylene imine, a polyamide having a general formula [-NH(CH₂)_r-NHCO(CH₂)_rCO-]_m, polyamides of
30 general formula [-NH(CH₂)_rCO-]_m, polyesters of general formula [-O(CH₂)_rCO-]_m, alkylated silyl compounds of general formula -Si(Y)₃, hetero-oligo/polyaminoacids of the general formula (-NHCHaaCO-)_m, a polyethylene glycol of the general formula -
35 (CH₂CH₂O)_m-CH₂CH₂OH, and a monoalkylated polyethylene glycol of the general formula (CH₂CH₂O)_m-CH₂CH₂OR.

hexyl, r is in the range of 1 to 20, and aa represents any amino acid side chain of naturally occurring amino acids.

25. The method according to claim 13, wherein the mass-modified nucleic acid is prepared from a single-stranded template polynucleotide complementary to the nucleic acid sequence by synthesizing the nucleic acid using mass-modified nucleotides.
26. The method according to claim 25, wherein the mass-modified nucleic acid is synthesized using a primer having a sequence which allows the nucleic acid to be anchored to the solid support.
27. The method according to claim 25, wherein the mass-modified nucleic acid is synthesized using a DNA polymerase and mass-modified deoxyribonucleoside triphosphates (dNTPs).
28. The method according to claim 25, wherein the mass-modified nucleic acid is synthesized using an RNA polymerase and mass modified ribonucleoside triphosphates (NTPs).
29. The method according to claim 25, wherein synthesis of the nucleic acid is initiated in the presence of an initiator oligonucleotide having a 5'-functionality allowing the nucleic acid to be immobilize on the solid support.
30. The method according to claim 9, wherein the nucleic acid further comprises a linking group (L) for covalently attaching the nucleic acid to the solid support.
31. The method according to claim 30, wherein the solid support further comprises a splint oligonucleotide and the linking group (L) comprises a nucleotide sequence able to anneal to the splint oligonucleotide and be covalently attached to the solid support by action of a ligase activity.

32. A system for exonuclease-mediated mass spectrometric sequencing comprising
- (i) reactor means for containing exonuclease cleavage reactions of the target nucleic acids, the cleavage reaction forming a train of sequentially released individual nucleotides;
 - (ii) mass spectrometric means for detecting the train of individual nucleotides released by the exonuclease cleavage of the target nucleic acid; and
 - (iii) transfer means for transferring the train of individual nucleotides from the reactor means to the mass spectrometric means.
33. The system of claim 32, wherein the transfer means further comprises a moving belt.
34. The system according to claim 32, wherein the reactor means further comprises a reactor with a cooling/heating mantle for controlling the temperature of the exonuclease cleavage reaction, at least one reagent flask for providing at least one reagent to the reactor, at least one pumping device for pumping the reagents and reactants to and from the reactor, at least one heating/cooling coil for controlling the temperature of reagents provided to the reactor, and means for immobilizing one of the target nucleic acids or the exonuclease in the reactor.
35. The system according to claim 34, wherein the reactor means further comprises at least one secondary reactor for mass-modifying the individual nucleotides released by the exonuclease cleavage of the target nucleic acids.
36. The system according to claim 35, wherein the secondary reactor comprises immobilized alkaline phosphatase.
37. The system according to claim 33, in which the moving belt comprises an endless moving belt driven by a stepping motor and held under tension by spring-loaded pulleys, a sample application area for applying a sample of the individual nucleotides released by the exonuclease cleavage of the target nucleic acids to a portion of the moving belt, a cooling/heating plate positioned beneath the sample application area for controlling the temperature of the sample application area, a CCD camera with optics for viewing the sample application area, a source for ionization/desorption capable of moving across the surface of the belt in a direction perpendicular to the motion of the moving belt, differential vacuum pumping stages for generating a vacuum over a portion of the moving belt, and a heating

38. The system according to claim 37, further comprising a washing station to wash the moving belt with a cleaning solution prior to application of the sample, wherein the heating device facilitates drying the belt after rinsing with the cleaning solution.
39. The system of claim 37, wherein the heating device is a microwave.
40. The system according to claim 37, in which the ionization/desorption source is a laser.
41. The system according to claim 32, further comprising a plurality of reactor means for parallel sequencing of a plurality of nucleic acids.
42. The system of claim 32, further comprising a microprocessor for processing the detected train of released nucleotides and determining the sequence of the target nucleic acids.
43. A kit for exonuclease sequencing at least two different species of nucleic acids by mass spectrometry, comprising
- (i) an exonuclease for cleaving the nucleic acids unilaterally from a first end to sequentially release individual nucleotides;
- (ii) a set of nucleotides for synthesizing the different species of nucleic acids, at least a portion of the nucleotides being mass-modified such that sequentially released nucleotides of each of the different species of nucleic acids are distinguishable,
- (iii) a polymerase for synthesizing the nucleic acids from complementary templates and the set of nucleotides, and
- (iv) a solid support for immobilizing one of the nucleic acids or the exonuclease.
44. The kit of according to claim 43, wherein the nucleic acids are 2'-deoxyribonucleic acids (DNA).
45. The kit of according to claim 43, wherein the nucleic acids are ribonucleic acids (RNA).
46. The kit of according to claim 43, wherein the exonuclease is selected from a group consisting of snake venom phosphodiesterase, spleen phosphodiesterase, Bal-31 nuclease, E. coli exonuclease I, E. coli exonuclease VII, Mung Bean Nuclease, S1 Nuclease, and T1 Nuclease.

exonuclease activity of T7 DNA polymerase, an exonuclease activity of Taq DNA polymerase, an exonuclease activity of Deep Vent DNA polymerase, and an exonuclease activity of Vent_r DNA polymerase.

- 5 47. The kit according to claim 43, wherein the solid support is selected from a group consisting of a capillary, a flat membrane, glass beads, cellulose beads, polystyrene beads, Sephadex beads, Sepharose beads, polyacrylamide beads and agarose beads.
48. The kit according to claim 43, wherein the solid support is functionalized to
10 facilitate covalent attachment of the nucleic acids.
49. The kit according to claim 43, wherein the exonuclease is immobilized by covalent attachment to the solid support.
- 15 50. The kit according to claim 43, wherein the mass-modified nucleotide comprises a mass-modifying functionality (M) attached to a nucleotide moiety.
51. The method according to claim 50, wherein the mass-modifying functionality (M) is attached to a nucleotide moiety at a position selected from a group consisting of a 5'
20 phosphate, a C-2' position of a sugar moiety, and a heterocyclic base.
52. The method according to claim 50, wherein the mass modifying functionality (M) is selected from a group consisting of XR, F, Cl, Br, I, Si(CH₃)₃, Si(CH₃)₂(C₂H₅), Si(CH₃)(C₂H₅)₂, Si(C₂H₅)₃, CH₂F, CHF₂, and CF₃, wherein X is selected from a group
25 consisting of -OH, -NH₂, -NHR, -SH, -NCS, -OCO(CH₂)_rCOOH (where r = 1-20), -NHCO(CH₂)_rCOOH (where r = 1-20), -OSO₂OH, -OCO(CH₂)_rI (where r = 1-20), and -OP(O-Alkyl)N(Alkyl)₂, and R is selected from a group consisting of H, methyl, ethyl, propyl, isopropyl, t-butyl, hexyl, benzyl, benzhydryl, trityl, substituted trityl, aryl, substituted aryl, polyoxymethylene, monoalkylated polyoxymethylene, a polyethylene imine, a
30 polyamide having a general formula [-NH(CH₂)_r NHCO(CH₂)_rCO-]_m, polyamides of general formula [-NH(CH₂)_r CO-]_m, polyesters of general formula [-O(CH₂)_rCO-]_m, alkylated silyl compounds of general formula -Si(Y)₃, hetero-oligo/polyaminoacids of the general formula (-NHCHaaCO-)_m, a polyethylene glycol of the general formula -
(CH₂CH₂O)_m-CH₂CH₂OH, and a monoalkylated polyethylene glycol of the general
35 formula -(CH₂CH₂O)_m-CH₂CH₂O-Y, where m is in the range of 0 to 200, Y is a lower alkyl group selected from a group consisting of methyl, ethyl, propyl, isopropyl, t-butyl, hexyl, r is in the range of 1 to 20, and aa represents any amino acid side chain of naturally

53. The kit according to claim 43, further comprising an instruction manual providing the endonuclease sequencing protocol.
54. A kit for sequencing nucleic acid by exonuclease-mediated mass spectrometry,
5 comprising
- (i) an exonuclease for cleaving the nucleic acid unilaterally from a first end to sequentially release individual nucleotides;
 - (ii) a set of nucleotides for synthesizing the nucleic acid, at least a portion of the nucleotides being mass-modified to modulate cleavage activity of the exonuclease,
 - 10 (iii) a polymerase for synthesizing the nucleic acids from complementary templates and the set of nucleotides, and
 - (iv) a solid support for immobilizing one of the nucleic acids or the exonuclease.
- 15 55. A method of determining a sequence of a nucleic acid, comprising
- (i) isolating the nucleic acid to be sequenced;
 - (ii) cleaving the nucleic acid unilaterally from a first end with an exonuclease activity to produce a set of nested nucleic acid fragments;
 - (iii) determining the molecular weight value of each one of the set of
 - 20 nucleic acid fragments by mass spectrometry; and
 - (iv) determining the sequence of the nucleic acid from the molecular weight values of the set of nucleic acid fragments.

FIG. 1





FIG. 2



DNA POL

FIG. 3




FIG. 4A






FIG. 4B



5 / 15

FIG. 5



$n = 1 - 1000$
 $M = H, OH, XR,$
 Halogen, N_3

FIG. 6













FIG. 7

Mass modifying functionality R (X similar to FIGURE 8):

H

Alkyl: $-(CH_2)_r-CH_3$ e.g. $-CH_3$, $-C_2H_5$,
and branched e.g. $-CH(CH_3)_2$

$ICH_2(CH_2)_r-O-H$

2,3-Epoxy-1-propanol

$-(CH_2)_m-CH_2-O-H$

$-(CH_2)_m-CH_2-O-Alkyl$

$-(CH_2CH_2NH)_m-CH_2CH_2-NH_2$

$-[NH-(CH_2)_r-NH-C(=O)-(CH_2)_r-C(=O)]_m-NH-(CH_2)_r-NH-C(=O)-(CH_2)_r-C(=O)-OH$

$-[NH(CH_2)_r-C(=O)]_m-NH-(CH_2)_r-C(=O)-OH$

$-[NH-CHY-C(=O)]_m-NH-CHY-C(=O)-OH$

$-[O-(CH_2)_r-C(=O)]_m-O-(CH_2)_r-C(=O)-OH$

-S-

-Si(Alkyl)₃

-Halogen

-N₃

$-CH_2F$, $-CHF_2$, $-CF_3$

$m = 0, 1 - 200$


$r = 1 - 20$

8 / 15

FIG. 8


x
-O-
-O-C-(CH ₂) _r C-O-
-NH-C-/-C-NH-
-NH-C-(CH ₂) _r -C-O-
-NH-C-NH-
-O-P-O-Alkyl
-O-SO ₂ -O-
-O-C-CH ₂ -S-
-S-
-NH-

9 / 15



10 / 15

FIG. 10



SUBSTITUTE SHEET (RULE 26)

11 / 15

FIG. II






FIG. 12




13 / 15

FIG. 13




14 / 15

FIG. 14A



15 / 15

FIG. 14B



INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/US 94/02938A. CLASSIFICATION OF SUBJECT MATTER
C 12 Q 1/68, G 01 N 33/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C 12 Q, G 01 N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, A1, 89/03 432 (UNITED STATES DEPARTMENT OF ENERGY) 20 April 1989 (20.04.89), abstract; claims. --	1, 4
A	WO, A1, 89/12 694 (GENOMYX, INC.) 28 December 1989 (28.12.89), abstract. --	1, 22- 24, 32
P, A	US, A, 5 288 644 (BEAVIS et al.) 22 February 1994 (22.02.94), abstract; claims. ----	1, 32, 55

☐ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
09 June 1994Date of mailing of the international search report
01 07

Mailing address of the ISA

European Patent Office
P.O. Box 2280, HV Ruiswijk
The Netherlands
Fax (+31-70) 340-2040, Telex 513996
Fax (+31-70) 340-3016

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/US 94/02938 SAE 87863

In diesem Anhang sind die Mitglieder
der Patentfamilien der im obenge-
nannten internationalen Recherchenbericht
angeführten Patentdokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visé ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WO A1 8903432	20-04-89	CA A1 1314247 EP A1 381693 EP A4 381693 IL A0 87925 IL A1 87925 JP T2 3502041 US A 4962037	09-03-93 16-08-90 19-08-92 31-03-89 18-06-93 16-05-91 09-10-90
WO A1 8912694	28-12-89	AU A1 38477/89 AU B2 632756 EP A1 379548 JP T2 3500093 US A 5003059 US A 5174962	12-01-90 14-01-93 01-08-90 10-01-91 26-03-91 29-12-92
US A 5288644	22-02-94	keine - none - rien	